

Gamma Rays from I^{128}

OSCH, R. RADICELLA,* P. REYES, AND T. URSTEIN
Nuclear Spectroscopy Laboratories, Comisión Nacional de Energía Atómica, Buenos Aires, Argentina
 (Received February 10, 1958)

The gamma radiation following the decay of I^{128} was studied by means of a single-channel analyzer. Four gamma rays of energies 0.445, 0.530, 0.740, and 0.975 Mev and relative intensities 100:9.3:0.9:1.8 were found. Three methods of preparing the source were employed in order to check the purity of the sample. Using published data together with our results, we propose a level sequence for Xe^{128} , which is of vibrational type.

I. INTRODUCTION

THE I^{128} nucleus disintegrates by K capture and β^- emission to two stable isobars, Te^{128} and Xe^{128} , respectively. A half-life of 24.99 min¹ was assigned to this disintegration. The K -capture branch is $6.3 \pm 0.7\%$ of that of electron branch.² According to Mims and Halban,² the x-rays appearing in the disintegration of I^{128} belong to the Te isotope.

Wapstra *et al.*³ observed two gamma rays of 0.445 and 0.980 Mev, Germagnoli⁴ one of 0.430 Mev. The β^- and γ radiations following the decay of I^{128} have been studied in detail by Benczer, Farrelly, Koerts, and Wu.⁵ They observed gamma rays of energies 0.455, 0.540, 0.750, and 0.990 Mev. Gupta and Jha⁶ have also studied the gamma-ray spectrum and observed 0.440- and 0.980-Mev gamma rays.

We continued the investigation of I^{128} because, as Xe^{128} is an even-even nucleus whose neutron number lies between 36 and 88, one of the vibrational bands according to Scharff-Goldhaber and Weneser⁷ and Wilets and Jean,⁸ there should exist at approximately twice the energy of the first excited state, three levels with character $0+$, $2+$, and $4+$. As the β^- transitions from I^{128} to the levels $0+$ and $2+$ of Xe^{128} are allowed (as will be seen later on), it is logical to expect the observation of the levels $0+$ and $2+$ of the triplet. This is of particular interest because one does not know with certainty the position of the $0+$ with respect to the $2+$ and $4+$ levels of the triplet.⁹ Recent theoretical work of Raz¹⁰ indicates that the $0+$ level lies higher in energy than the $2+$ and $4+$ levels and that the $2+$ level is always below the $4+$. There is only one example in which the three sublevels appear¹¹ (Cd^{114})

* Radiochemical Laboratories.

¹ D. Hull and H. Seeling, *Phys. Rev.* **60**, 553 (1941).

² J. Reynolds, *Phys. Rev.* **79**, 745 (1950); H. Mims and H. Halban, *Proc. Phys. Soc. (London)* **A64**, 753 (1953).

³ H. Wapstra *et al.*, *Physica* **19**, 138 (1953).

⁴ E. Germagnoli *et al.*, *Nuovo cimento* **10**, 1388 (1953).

⁵ Benczer, Farrelly, Koerts, and Wu, *Phys. Rev.* **101**, 1027 (1956).

⁶ R. K. Gupta and S. Jha, *Nuclear Phys. I*, **2** (1956).

⁷ G. Scharff-Goldhaber and J. Weneser, *Phys. Rev.* **98**, 212 (1955).

⁸ L. Wilets and M. Jean, *Phys. Rev.* **102**, 788 (1956).

⁹ C. A. Mallmann (to be published).

¹⁰ B. J. Raz (to be published).

¹¹ H. T. Motz, *Phys. Rev.* **104**, 1352 (1956).

and one in which the sequence $2+$, $0+$ was found¹² (Pd^{106}). In the latter case, the energy of the corresponding two levels is the same within the experimental errors.

II. EXPERIMENTAL

(A) Source Preparation

Different methods of preparing the source were employed in order to check the purity of the sample. I^{128} was prepared by bombarding ethyl iodide with slow neutrons from the synchrocyclotron of this Institute, and the iodide separated by Szilard-Chalmers method. I^{128} was also prepared by bombarding re-sublimed iodide with slow neutrons. Finally, NaI was irradiated with slow neutrons and the iodine separated. A half-life determination for the total β^- activity was performed with a scintillation spectrometer with anthracene in the three cases, and the same results were obtained. In Fig. 1 the decay curve of the β^- activity is plotted.

(B) Single Measurements

For the measurements of energies and relative intensities of the gamma radiations, a single scintillation spectrometer with crystals of different sizes was used.




FIG. 1. Decay curve of the β^- activity corresponding to I^{128} .

¹² D. E. Alburger and B. J. Toppel, *Phys. Rev.* **100**, 1457 (1955).

Final measurements were made with a 2-in. \times 2-in. NaI(Tl) crystal and a DuMont 6364 photomultiplier. Many observations at different source-to-crystal distances were performed in order to estimate the sumup between the gamma rays.

A typical gamma-ray spectrum of I^{128} after correction for decay is plotted in Fig. 2. In the insert is represented the spectrum for low energies. The following energies, in Mev, were established: 0.0275 (Te x-rays), 0.445 ± 0.005 , 0.530 ± 0.005 , 0.740 ± 0.010 , and 0.975 ± 0.015 .

In order to reduce the pileup to a few percent between the 0.445-Mev gamma ray and its Compton background and the sumup between the 0.445- and 0.530-Mev gamma rays, the source-to-crystal distance was fixed at 13 cm. The relative intensities of the gamma rays are listed in Table I.

III. DISCUSSION

It is possible to conclude from the series of measurements that there are four gamma rays following the




Fig. 2. Gamma-ray spectrum of I^{128} after correction for decay. In the insert is represented the spectrum for low energies.

decay of I^{128} whose energies are, in Mev: 0.0275 (Te x-rays), 0.445 ± 0.005 , 0.530 ± 0.005 , 0.740 ± 0.010 , and 0.975 ± 0.015 .

The ground state of even-even nuclei always has the character $0+$.^{9,10} The first excited state, with few exceptions, has the character $2+$.⁹⁻¹¹ The energy of the first excited level was plotted as a function of neutron number for a series of Te and Xe nuclei (Fig. 3). It appears reasonable to compare these energies, as the first excited states of all these nuclei have the character $2+$. This is not the case with the second excited states where the character is generally $2+$ or $4+$ and occasionally $0+$ or odd.^{9,11}

From the results of Temmer and Heydenburg¹³ and the plot of Fig. 3, we may assert that the 0.740-Mev gamma ray corresponds to the Te branch. According to that plot, the 0.445-Mev gamma ray is the corre-

TABLE I. Relative intensities of the gamma radiations.

Energy of the gamma radiation (Mev)	Relative intensities
0.445	100
0.530	9.3
0.740	0.9
0.975	1.8

sponding transition between the first and ground states of Xe^{128} .

According to the energy and relative intensities of the β^- radiations from I^{128} inferred by other authors,⁵ all the β^- transitions are allowed. As was stated before, the character of the ground and first excited levels of Xe^{128} are $0+$ and $2+$, respectively; consequently, the character of the ground state of I^{128} should be $1+$. This can be explained by the shell model: we can assign a configuration d_5 for protons and d_3 for neutrons. Then the character of the second excited level of Xe^{128} could be $0+$, $1+$, or $2+$.

In general the second excited levels (collective levels) of even-even nuclei have the character $2+$ or $4+$.⁸⁻¹⁰ The character $1+$ for the second excited state of even-even nuclei has not been observed. This was pointed out by Glaubman and Morinaga¹⁴ who asserted that in even-even nuclei the low-lying odd-parity states have odd spin. With this assumption we can exclude the possibility $1+$ for the second excited level of Xe^{128} . Owing to the fact that the energy of the 0.975-Mev gamma ray is 2.2 times that of the 0.445-Mev gamma ray and is the sum of 0.445 and 0.530 Mev, we are tempted to consider that radiation as the crossover from the second excited level to the ground state, and the possibility $0+$ must be ruled out.

We did not observe any splitting in the second excited state of Xe^{128} , even though the $0+$ sublevel could be fed by an allowed β^- transition. In order to estimate the energy and relative intensity of the possible transition from this sublevel to the $2+$ first excited state, we have made a rough calculation, obtaining the following results: If we suppose a β^- transition to the




Fig. 3. Energy of the first excited level of a series of Te and Xe nuclei, as a function of neutron number.

¹³ G. M. Temmer and N. P. Heydenburg, Phys. Rev. **104**, 967 (1956).

¹⁴ M. J. Glaubman, Phys. Rev. **90**, 1000 (1953); H. Morinaga, Phys. Rev. **103**, 503 (1956).

$0+$ second excited sublevel with a $\log ft$ value from 6.5 to 7 and a gamma transition of intensity at most 10% of that of the 0.445-Mev gamma ray, its corresponding energy would lie between 0.230 and 0.440 Mev. On the other hand, if we suppose that the gamma transition has an intensity of 1% or less of that of the 0.445-Mev gamma ray, its corresponding energy would lie between 0.530 and 0.740 Mev. These rough calculations are not in contradiction with our experimental results, since these relative intensities at the corresponding energy ranges cannot be observed.

We shall attempt to perform the external conversion of the gamma rays from I^{128} and observe the corresponding electron lines in our orange-type beta-ray spectrometer.

ACKNOWLEDGMENTS

We are very much indebted to Dr. C. A. Mallmann for his valuable discussions throughout this work, especially for his advice about the systematics of even-even nuclei. We wish also to thank the synchrocyclotron group for their cooperation.